



OPEN ACCESS



Operative treatment versus nonoperative treatment of Achilles tendon ruptures: systematic review and meta-analysis

Yassine Ochen,^{1,2} Reinier B Beks,^{1,3} Mark van Heijl,^{3,4} Falco Hietbrink,¹ Luke P H Leenen,¹ Detlef van der Velde,⁵ Marilyn Heng,² Olivier van der Meijden,⁶ Rolf H H Groenwold,⁷ R Marijn Houwert¹

For numbered affiliations see end of article.

Correspondence to:

Yassine Ochen
yassineochen@gmail.com

(ORCID 0000-0002-7187-2055)

Additional material is published online only. To view please visit the journal online.

Cite this as: *BMJ* 2018;364:k5120
<http://dx.doi.org/10.1136/bmj.k5120>

Accepted: 14 November 2018

ABSTRACT

OBJECTIVES

To compare re-rupture rate, complication rate, and functional outcome after operative versus nonoperative treatment of Achilles tendon ruptures; to compare re-rupture rate after early and late full weight bearing; to evaluate re-rupture rate after functional rehabilitation with early range of motion; and to compare effect estimates from randomised controlled trials and observational studies.

DESIGN

Systematic review and meta-analysis.

DATA SOURCES

PubMed/Medline, Embase, CENTRAL, and CINAHL databases were last searched on 25 April 2018 for studies comparing operative versus nonoperative treatment of Achilles tendon ruptures.

STUDY SELECTION CRITERIA

Randomised controlled trials and observational studies reporting on comparison of operative versus nonoperative treatment of acute Achilles tendon ruptures.

DATA EXTRACTION

Data extraction was performed independently in pairs, by four reviewers, with the use of a predefined data extraction file. Outcomes were pooled using random effects models and presented as risk difference, risk ratio, or mean difference, with 95% confidence interval.

RESULTS

29 studies were included—10 randomised controlled trials and 19 observational studies.

The 10 trials included 944 (6%) patients, and the 19 observational studies included 14 918 (94%) patients. A significant reduction in re-ruptures was seen after operative treatment (2.3%) compared with nonoperative treatment (3.9%) (risk difference 1.6%; risk ratio 0.43, 95% confidence interval 0.31 to 0.60; $P<0.001$; $I^2=22\%$). Operative treatment resulted in a significantly higher complication rate than nonoperative treatment (4.9% v 1.6%; risk difference 3.3%; risk ratio 2.76, 1.84 to 4.13; $P<0.001$; $I^2=45\%$). The main difference in complication rate was attributable to the incidence of infection (2.8%) in the operative group. A similar reduction in re-rupture rate in favour of operative treatment was seen after both early and late full weight bearing. No significant difference in re-rupture rate was seen between operative and nonoperative treatment in studies that used accelerated functional rehabilitation with early range of motion (risk ratio 0.60, 0.26 to 1.37; $P=0.23$; $I^2=0\%$). No difference in effect estimates was seen between randomised controlled trials and observational studies.

CONCLUSIONS

This meta-analysis shows that operative treatment of Achilles tendon ruptures reduces the risk of re-rupture compared with nonoperative treatment. However, re-rupture rates are low and differences between treatment groups are small (risk difference 1.6%). Operative treatment results in a higher risk of other complications (risk difference 3.3%). The final decision on the management of acute Achilles tendon ruptures should be based on patient specific factors and shared decision making. This review emphasises the potential benefits of adding high quality observational studies in meta-analyses for the evaluation of objective outcome measures after surgical treatment.

Introduction

Rupture of the Achilles tendon is a frequently encountered injury, with an incidence of 31 per 100 000 per year, and is most common in the young to middle aged active population, with a reported mean age ranging from 37 to 44 years.^{1,2} Recent studies indicate that the incidence of Achilles tendon rupture is still increasing owing to a more active older population.² Injury of the Achilles tendon can be debilitating because of its role in ambulation and activity, affecting both athletes and non-athletes. The management of acute Achilles tendon ruptures—operative or nonoperative treatment—is much debated.²

WHAT IS ALREADY KNOWN ON THIS TOPIC

Rupture of the Achilles tendon is a frequently encountered injury, with recent studies indicating an increase in incidence of Achilles tendon ruptures. Meta-analyses of randomised controlled trials have shown operative treatment to significantly reduce the risk of tendon re-rupture compared with nonoperative treatment (reported risk difference 5-7%). However, operative treatment leads to a significant increase in other complications compared with nonoperative treatment, with a reported risk difference of 16-21%.

WHAT THIS STUDY ADDS

Operative treatment of acute Achilles tendon ruptures reduced the risk of re-rupture compared with nonoperative treatment. However, re-rupture rates were low and differences between treatment groups were small, with a risk difference of 1.6%. Operative treatment resulted in a higher risk of other complications than nonoperative treatment (risk difference 3.3%), mostly attributable to increased risk of infection.

Several meta-analyses of randomised controlled trials (RCTs) have shown that operative treatment significantly reduces the risk of tendon re-rupture compared with nonoperative treatment, with a reported risk difference in re-rupture rate varying from 5% to 7%.³⁻⁶ However, operative treatment leads to a significant increase in other complications such as infection, deep vein thrombosis, and sural nerve injury, with a reported risk difference varying from 16% to 21%.³⁻⁶ The incidence of operative treatment has declined over the past decade as a result of multiple RCTs showing comparable results between operative and nonoperative treatment.^{1,2}

A recent systematic review of overlapping meta-analyses evaluated nine meta-analyses that compared operative and nonoperative treatment of Achilles tendon ruptures. The discordance found among the nine meta-analyses indicated that further investigation is warranted as rehabilitation protocols, weight bearing restrictions, and treatment modalities have evolved.⁷

Systematic reviews and meta-analyses of RCTs are considered the highest level of evidence for the evaluation of treatment effects. However, several reports have shown that little evidence exists for significant differences in effect estimates between RCTs and observational studies.⁸⁻¹¹ The addition of observational studies in meta-analyses increases sample size, which could enable the evaluation of small treatment effects and infrequent outcome measures. Furthermore, observational studies might provide insight into a variety of populations and long term effects compared with the usually highly selected patient populations in RCTs.¹²⁻¹³ Both RCTs and observational studies are increasingly used in orthopaedic trauma meta-analyses for the evaluation of treatment effects.¹⁴⁻¹⁷

The primary aim of this systematic review and meta-analysis was to compare re-rupture rate, complication rate, and functional outcome after operative versus nonoperative treatment of acute Achilles tendon ruptures. Secondly, we sought to evaluate re-rupture rate after early and late full weight bearing and compare re-rupture rate after functional rehabilitation with early range of motion. Finally, we compared effect estimates obtained from RCTs and observational studies.

Methods

This systematic review and meta-analysis was performed and reported according to the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) and the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklists.¹⁸⁻²⁰ A published protocol for this review does not exist.

Search strategy and selection criteria

We last searched the PubMed/Medline, Embase, CENTRAL, and CINAHL databases on 25 April 2018 for studies comparing operative versus nonoperative treatment of Achilles tendon ruptures. The search

syntax is provided in supplementary table A. Duplicate articles were removed. Two reviewers (YO, RHHG) independently screened titles and abstracts for eligibility of identified studies. All published comparative studies, both RCTs and observational studies, reporting on the comparison of operative versus nonoperative treatment of acute Achilles tendon ruptures were eligible for inclusion.

After title and abstract screening, the same two reviewers (YO, RHHG) independently reviewed full text articles. Inclusion criteria were acute Achilles tendon rupture, operative treatment (open or minimally invasive surgery) versus nonoperative treatment (cast immobilisation or functional bracing), treatment within four weeks of rupture, age 16 years or older, and reporting of re-rupture rate, complication rate, or functional outcome. Exclusion criteria were delayed presentation (treatment more than four weeks after rupture), treatment for re-rupture, language other than English, no availability of full text article, and letters, meeting proceedings, and case reports. We had no inclusion restrictions based on weight bearing status or functional rehabilitation protocol. Disagreements on eligibility of full text articles were resolved by consensus or by discussion with a third reviewer (RMH). References of included studies were screened, and backwards citation tracking was performed using Web of Science to identify articles not found in the original literature search.

Data extraction

Four reviewers (YO, RHHG, RMH, RBB) extracted data independently in pairs, using a predefined data extraction file. The following baseline characteristics were extracted from the included studies: first author, year of publication, study design, country in which the study was performed, study period, number of included patients, operative method, nonoperative method, full weight bearing status, functional rehabilitation protocol, and mean follow-up. Studies reporting on patient cohorts described in previously published articles were excluded or merged.

Quality assessment

The same four reviewers (YO, RHHG, RMH, RBB), in pairs, independently assessed the methodological quality of included studies by using the Methodological Index for Non-Randomised Studies (MINORS).²¹ The MINORS is a validated instrument for the assessment of methodological quality and clear reporting of non-randomised surgical studies, resulting in a score ranging from 0 to 24 for comparative studies.²¹ In this study the assessment of methodological quality resulted in a score ranging from 0 to 24 for RCTs and prospective cohort studies. The methodological quality of retrospective cohort studies resulted in a score ranging from 0 to 18. The MINORS criteria for prospective collection of data, loss to follow-up, and prospective calculation of study size were not applicable to the retrospective cohort studies. Details on the methodological quality assessment are

provided in supplementary table B. Disagreements were resolved by consensus.

Primary and secondary outcomes

The primary outcome measure was re-rupture rate after operative or nonoperative treatment. Secondary outcome measures included complication rate, functional outcome scores, return to sporting activity, and return to work after operative or nonoperative treatment. We defined complication rate as the rate of complications other than re-rupture. Complications included reports of wound infection, sural nerve injury, deep vein thrombosis, and pulmonary embolism. Functional outcome scores included the Achilles Tendon Rupture Score (ATRS).²² We subdivided functional outcome scores according to follow-up, into short term (one year or less) and long term (more than one year). We defined return to sporting activity as the duration in months before resumption of sports and return to work as the duration in weeks before resuming work. In studies that reported on both open and minimally invasive surgery, we used the combined outcome measures.

Statistical analysis

We present all continuous variables as mean value with standard deviation or range. We converted continuous variables to mean and standard deviation if sufficient information was available, using the methods described in the Cochrane Handbook for Systematic Reviews of Interventions.²³ We extracted dichotomous variables as absolute number and percentage, pooled them using the Mantel-Haenszel method, and presented them as risk difference and risk ratio with 95% confidence interval. We pooled continuous outcomes by using the inverse variance weighting method and presented them as mean difference with 95% confidence interval. We used random effects models for all analyses. We assessed statistical heterogeneity between studies by visual inspection of forest plots and by the I^2 and χ^2 statistics for heterogeneity. We used the overall effect Z test to determine the significance level for treatment effects. All analyses were stratified according to study design—RCTs or observational studies. We assessed difference in effect estimates between the two subgroups as described in the Cochrane Handbook for Systematic Reviews of Interventions.²³ The significance level for difference in effect estimates across the subgroups was determined by the test for subgroup differences. We defined the significance level for treatment effects and differences across the subgroups as a P value below 0.05. We assessed potential publication bias by visual inspection of funnel plots with risk ratio and standard error.²⁴ We used Review Manager (RevMan, version 5.3.5) for all statistical analyses.²⁵ We further assessed publication bias with Begg's and Egger's statistical tests using Stata 13.1.

Primary sensitivity analyses

We did sensitivity analyses for the primary outcome, including studies with an early (four weeks or less)

and late (more than four weeks) full weight bearing status after treatment. Studies reporting on both an early and a late full weight bearing cohort were accordingly divided for sensitivity analysis. We did an additional sensitivity analysis for the primary outcome with studies that included an accelerated functional rehabilitation protocol. We defined accelerated functional rehabilitation as the start of early range of motion within three weeks after nonoperative treatment. Rehabilitation with functional bracing systems with successive fixed degrees of plantar flexion, which did not allow for free range of motion, were not considered as accelerated rehabilitation.

Secondary sensitivity analyses

We did secondary sensitivity analyses for high quality studies and year of study period, regarding re-rupture rate and complication rate. We defined high quality studies as RCTs or prospective cohort studies with a MINORS score of 16 or higher (range 0-24) or retrospective cohort studies with a MINORS score of 12 or higher (range 0-18). We did additional sensitivity analyses with studies that included patients after the study period 2000, to account for the development of new rehabilitation protocols, operative techniques, and nonoperative treatment modalities.

Patient and public involvement

No patients were involved in setting the research question or the outcome measures, nor were they involved in developing plans for or implementation of the study. No patients were asked to advise on interpretation or writing up of results. There are no plans to disseminate the results of the research to study participants or the relevant patient community.

Results

Search

Figure 1 shows a flowchart of the literature search and study selection. Full text articles could not be obtained for three studies.²⁶⁻²⁸ Four studies reported on patient cohorts described in previously published articles and were excluded or merged with the original studies.²⁹⁻³² This resulted in the final inclusion of 29 studies for analyses in this systematic review and meta-analysis—10 RCTs and 19 observational studies.³³⁻⁶¹

Baseline study characteristics

The 29 studies included 15 862 patients, of whom 9375 were treated operatively and 6487 nonoperatively. The overall weighted mean age was 41 (range 17-86) years, 41 years in the operative group and 44 years in the nonoperative group. Overall, the studies included 11 779 (74%) males. Overall follow-up ranged from 10 to 95 months. Table 1 shows the baseline characteristics for both RCTs and observational studies. In addition, supplementary table C shows the treatment characteristics of all included studies.

The 10 RCTs included 944 (6%) patients; 469 patients were treated operatively and 475




Fig 1 | PRISMA flow diagram representing search and selection of studies comparing operative versus nonoperative treatment of Achilles tendon ruptures

nonoperatively. The weighted mean age was 40 years in both treatment groups, and 779 (83%) males were included. The operative method was open surgery in nine studies and minimally invasive surgery in one study.

The 19 observational studies—three prospective and 16 retrospective cohort studies—included 14 918 (94%) patients. Operative treatment was performed in 8906 patients, and 6012 were treated nonoperatively. The weighted mean age in the studies was 42 (range 17-86) years, 40 years in the operative group and 44 years in the nonoperative group, and 11 000 (74%) patients were male. The operative method was open

surgery in nine studies, unclear in four studies, and a combination of open and minimally invasive surgery in six studies.

Quality assessment

The overall mean MINORS score was 14.3 (SD 5.2; range 5-23). The mean MINORS score for the RCTs was 20.3 (2.6; 16-23). The mean MINORS score for the observational studies was 11.2 (2.8; 5-16), 14 (2; 12-16) for the prospective cohort studies and 10.6 (2.6; 5-15) for the retrospective cohort studies. The details and distribution of MINORS scores are provided in supplementary table D.

Table 1 | Baseline characteristics of studies included in meta-analysis of Achilles tendon ruptures

Study and year	Study period	Country	Overall No	Number		Mean (SD or range) age, years		Sex (female/male)		Mean (SD or range) follow-up, months	
				OP	NON	OP	NON	OP	NON	OP	NON
Randomised controlled trials											
Cetti et al, 1993	1982-84	Denmark	111	56	55	37.2 (21-62)	37.8 (21-65)	9/47	10/45	12	12
Keating et al, 2011	2000-04	UK	80	39	41	41.2 (27-59)	39.5 (21-58)	11/28	9/32	12	12
Lantto et al, 2016	2009-13	Finland	60	32	28	40 (27-57)	39 (28-60)	2/30	3/25	18	18
Metz et al, 2008	2004-05	Netherlands	83	42	41	40 (23-63)	41 (25-62)	11/31	6/35	12	12
Möller et al, 2001	1995-97	Sweden	112	59	53	39.6 (21-63)	38.5 (26-59)	8/51	5/48	24	24
Nilsson-Helander et al, 2010	2004-07	Sweden	97	49	48	40.9 (8.8)	41.2 (9.5)	9/40	9/39	12	12
Nistor et al, 1981	1973-77	Sweden	107	46	61	Overall 41 (21-77)		Overall 11/96		Overall 30 (12-60)	
Olsson et al, 2013	2009-10	Sweden	100	49	51	39.8 (8.9)	39.5 (9.7)	10/39	4/47	12	12
Twaddle et al, 2007	1997-2002	New Zealand	50	25	25	41.8	40.3	6/14*	8/14*	12	12
Willits et al, 2010	2000-05	Canada	144	72	72	39.7 (11)	41.1 (8.0)	13/59	13/59	24	24
Observational studies											
Bergkvist et al, 2012	2002-06	Sweden	487	220	267	43 (11)	47 (14)	Overall 78/409		Overall 43 (12-97)	
Carden et al, 1987	1969-81	UK	71	35	36	42.7 (26-68)	43 (22-70)	10/26	12/25	48 (12-204)	64 (12-120)
Costa et al, 2006†	2001-02	UK	96	48	48	42 (28-69)	53 (21-79)	7/40*	16/32	12	12
Cukelj et al, 2015	1998-2013	Croatia	90	60	30	34.8 (4.7)	35.1 (4.7)	9/51	9/21	Overall 12	
Ebinesan et al, 2008	2001-03	UK	63	51	12	44.8	52.1	14/37	6/6	NA	
Fahlström et al, 1998	1990-94	Sweden	31	22	9	34.6 (23-50)	39.4 (28-51)	Overall 4/27		Overall 39 (16-67)	
Grubor et al, 2012	2003-10	Bosnia	42	34	8	NA	NA	Overall 5/37		Overall 12	
Gwynne-Jones et al, 2011	1999-2008	New Zealand	363	143	220	37.4	40.9	59/84	107/113	NA	
Jaakkola et al, 2001	1985-99	USA	73	35	38	37.3 (25-64)	38.0 (21-62)	3/32	6/32	43	54
Jackson et al, 2013†	2002-08	UK	80	29	51	37 (24-55)	47 (27-80)	3/26	16/35	NA	
Kotnis et al, 2006†	2000-05	UK	125	67	58	41.0 (26-80)	43.9 (26-85)	19/48	18/40	12	12
Lim et al, 2017	NA	New Zealand	200	99	101	40.1	42	21/41*	32/38*	Overall 78 (24-156)	
Miller et al, 2005	1990-96	UK	172	140	32	45	49	23/117	11/21	Overall 95 (53)	
Nestorson et al, 2000	1992-97	Sweden	24	14	10	72 (65-79)	71 (65-86)	3/11	1/9	Overall 39 (13-65)	
Rajasekar et al, 2005	1997-2001	UK	35	21	14	NA	NA	Overall 10/25		Overall 24 (9-48)	
Renninger et al, 2016	2011-14	USA	57	27	30	32.3 (25-40)	29.7 (23-44)	0/27	0/30	Overall 10	
Van der Linden et al, 2004	1990-2001	Netherlands	292	212	80	37 (9.4)	42 (12)	58/154	21/59	Overall 72 (36)	
Wang et al, 2015	2007-11	USA	12 570	7625	4945	NA	NA	1514/6111	1737/3208	NA	
Weber et al, 2003	1993-98	Switzerland	47	24	23	38 (28-51)	39 (17-55)	4/13*	8/15	49 (30-79)	23 (12-42)

NA=not available; NON=nonoperative treatment; OP=operative treatment.

*Ratio may not add up to total number of patients owing to loss to follow-up.

†Prospective cohort study; all other observational studies are retrospective cohort studies.

Primary outcome measure

Re-rupture rate

Re-rupture rate was reported in all 29 studies. The overall pooled effect showed that operative treatment was associated with a significant reduction in re-rupture rate compared with nonoperative treatment (risk ratio 0.43, 95% confidence interval 0.31 to 0.60; $P<0.001$; $I^2=22\%$) (fig 2). The pooled effect of RCTs showed a risk ratio of 0.40 (0.24 to 0.69; $P<0.001$; $I^2=0\%$). The pooled effect of observational studies showed a risk ratio of 0.42 (0.28 to 0.64; $P<0.001$; $I^2=31\%$). Re-rupture occurred in 2.3% of patients after operative treatment compared with 3.9% after nonoperative treatment (risk difference 1.6%). We found no significant difference in effect estimates from RCTs and observational studies (test for subgroup differences: $P=0.91$; $I^2=0\%$). There was no visual asymmetry in the funnel plot (supplementary figure A).

The Begg rank correlation test ($P=0.66$) and Egger linear regression test ($P=0.16$) indicated no evidence of publication bias.

Secondary outcome measures

Complication rate

Complication rate was reported in 26 (90%) studies—10 RCTs and 16 observational studies. The overall pooled effect showed a risk ratio of 2.76 (1.84 to 4.13; $P<0.001$; $I^2=45\%$) in favour of nonoperative treatment compared with operative treatment (fig 3). The pooled effect of RCTs showed a risk ratio of 3.26 (1.26 to 8.41; $P=0.01$; $I^2=74\%$). The pooled effect of observational studies showed a risk ratio of 2.93 (2.28 to 3.75; $P<0.001$; $I^2=0\%$). The incidence of complications was 4.9% after operative treatment compared with 1.6% after nonoperative treatment (risk difference 3.3%). Table 2 shows the classification and incidence of




Fig 2 | Forest plot of re-rupture rate in meta-analysis of Achilles tendon ruptures. M-H=Mantel-Haenszel

complications. The main complication after operative treatment was infection, which occurred in 2.8% of patients. The main complication after nonoperative treatment was deep vein thrombosis, which occurred in 1.2% of patients compared with 1.0% after operative treatment. We found no significant difference between effect estimates from RCTs and observational studies (test for subgroup differences: $P=0.83$; $I^2=0\%$).

There was no visual asymmetry in the funnel plot (supplementary figure B). The Begg rank correlation test ($P=0.50$) and Egger linear regression test ($P=0.11$) indicated no evidence of publication bias.

Functional outcome

Short term functional outcome assessed according to the ATRS score was reported in three (10%) studies.

BMJ: first published as 10.1136/bmj.k5120 on 7 January 2019. Downloaded from <http://www.bmj.com/> on 20 April 2024 by guest. Protected by copyright.

Nilsson-Helander et al reported a median ATRS score of 75 (range 31-100) in the operative group and 90 (31-100) in the nonoperative group.⁵⁸ Olsson et al reported a mean ATRS score of 82 (SD 20) in the operative group compared with 80 (23) in the nonoperative group.⁵⁹ In both RCTs, the differences found were non-significant. The observational study by Jackson et al reported a statistical significant difference in median ATRS score—94 (range 23-100) in the operative group and 84 (25-100) in the nonoperative group.³⁴

Long term functional outcome using the ATRS score was assessed in two observational studies. Bergkvist et al reported a mean ATRS score of 83 (SD 19) in the operative group and 78 (22) in the nonoperative group.³⁶ Lim et al reported a mean ATRS score of 85 in both groups.⁴⁴ No significant difference was found in either study. We did not pool functional outcome data owing to a wide variety in ATRS score reports and insufficient information to convert data. Descriptive details on functional outcome measures are provided in supplementary table E.




Fig 3 | Forest plot of complication rate in meta-analysis of Achilles tendon ruptures. M-H=Mantel-Haenszel

Table 2 | Number and incidence of complications in studies included in meta-analysis of Achilles tendon ruptures

Complication classification	Operative treatment		Nonoperative treatment	
	No	Incidence (%)	No	Incidence (%)
Pulmonary embolism	2	0.02	2	0.03
Deep vein thrombosis	89	0.97	74	1.17
Wound/skin infection	258	2.80	1	0.02
Sural nerve injury	39	0.42	5	0.08
Chronic pain	3	0.03	2	0.03
Scar/skin adhesion	35	0.38	15	0.24
Wound dehiscence	8	0.09	0	0
Not specified/other	21	0.23	3	0.05
Total	455	4.94	102	1.61

Return to sports and work

Return to sports was reported by four (14%) studies—one RCT and three observational studies (supplementary table E). The mean time varied between six and nine months after operative treatment and between six and eight months after nonoperative treatment. We could not pool data on return to sports in a meta-analysis, as only one study reported a mean and standard deviation.

Return to work was reported in nine (31%) studies—four RCTs and five observational studies (supplementary table E). The outcome data of six studies could not be pooled owing to insufficient reporting of information. The pooled effect estimates of three studies—two RCTs and one observational study—showed no significant mean difference between operative and nonoperative treatment groups (supplementary figure C).

Primary sensitivity analysis

Weight bearing status

Early (four weeks or less) weight bearing status was reported in nine (31%) studies—five RCTs and four observational studies. The overall pooled effect showed a significant reduction in re-rupture rate after operative treatment compared with nonoperative treatment in the early (four weeks or less) full weight bearing studies (risk ratio 0.49, 0.26 to 0.93; $P=0.03$; $I^2=9\%$) (supplementary figure D). Late (more than four weeks) weight bearing status was reported in 15 (52%) studies—four RCTs and 11 observational studies. The overall pooled effect of the late (more than four weeks) full weight bearing studies also showed a significant reduction in re-rupture rate in favour of operative treatment (risk ratio 0.33, 0.21 to 0.50; $P<0.001$; $I^2=0\%$) (supplementary figure E).

Accelerated functional rehabilitation

Accelerated functional rehabilitation with early range of motion was performed in six (21%) studies—three RCTs and three observational studies. The overall pooled effect showed no significant difference between operative and nonoperative treatment regarding re-rupture rate (risk ratio 0.60, 0.26 to 1.37; $P=0.23$; $I^2=0\%$) (fig 4).

Secondary sensitivity analyses

Table 3 shows the results of the secondary sensitivity analyses. Re-rupture rate was reported in 17 (59%)

high quality studies—10 RCTs and seven observational studies. The overall pooled effect showed that operative treatment was associated with a significant reduction in re-rupture rate compared with nonoperative treatment (risk difference 5.1%; risk ratio 0.44, 0.30 to 0.64; $P<0.001$; $I^2=0\%$) (supplementary figure F). Re-rupture rate was reported in 14 studies (48%) with a study period after the year 2000—six RCTs and eight observational studies. The overall pooled effect showed a significant reduction in re-rupture rate after operative treatment compared with nonoperative treatment (risk difference 0.9%; risk ratio 0.59, 0.42 to 0.83; $P=0.002$; $I^2=10\%$) (supplementary figure G).

Complication rate was reported in 16 (55%) high quality studies—10 RCTs and six observational studies. The overall pooled effect showed a risk ratio of 2.72 (1.44 to 5.12; $P=0.002$; $I^2=62\%$) in favour of nonoperative treatment compared with operative treatment (risk difference 8.8%) (supplementary figure H). Complication rate was reported in 14 (48%) studies with a study period after the year 2000—six RCTs and eight observational studies. The overall pooled effect showed a risk ratio of 2.15 (1.28 to 3.60; $P=0.004$; $I^2=52\%$) in favour of nonoperative treatment compared with operative treatment (risk difference 2.4%) (supplementary figure I).

Discussion

This systematic review and meta-analysis, including both RCTs and observational studies, compared outcomes after operative versus nonoperative treatment of acute Achilles tendon ruptures. The pooled effect estimate showed that operative treatment was associated with a significant reduction in re-rupture rate compared with nonoperative treatment. However, operative treatment resulted in a significantly higher rate of other complications. Sensitivity analyses showed a similar reduction in re-rupture rate after both early and late full weight bearing in favour of operative treatment compared with nonoperative treatment. However, we found no significant difference in re-rupture rate if accelerated functional rehabilitation with early range of motion was used. Sensitivity analyses with high quality studies and studies with a study period after the year 2000 also showed operative treatment to be associated with a significant reduction in re-rupture rate but a higher risk of other complications. We found no significant difference in




Fig 4 | Forest plot of re-rupture rate in studies that included accelerated functional rehabilitation in meta-analysis of Achilles tendon ruptures. M-H=Mantel-Haenszel

effect estimates from RCTs and observational studies, for either re-rupture rate or complication rate.

Comparison with previous findings

Operative treatment reduces the risk of re-rupture compared with nonoperative treatment, but it also results in a higher risk of other complications. These findings are in accordance with those of previous meta-analyses.³⁻⁶ Our review included 10 RCTs with a total of 944 patients, which resulted in an increased number of patients available for analyses, thus exceeding previous meta-analyses. Furthermore, the inclusion of observational studies resulted in an additional 14918 patients for analyses. The previous meta-analyses reported a risk difference in re-rupture rate varying from 5% to 7% and a risk difference of other complications varying from 16% to 21%.³⁻⁶ However, with the addition of observational studies, this review shows that differences between treatment groups are small, with a risk difference in re-rupture rate of 1.6% and a risk difference of 3.3% for other complications.

Functional outcome measures included the ATRS score, return to sports, and return to work. The ATRS score is the most commonly used patient reported

instrument to evaluate limitations after treatment for an acute Achilles tendon rupture.² ATRS scores were not pooled in this study, but most studies showed no significant difference in ATRS score between the operative and nonoperative treatment groups. Resumption of sports was reported by only four studies; the results indicate no difference between operative treatment (six to nine months) and nonoperative treatment (six to eight months). The pooled effect of return to work showed no significant difference between treatment groups. Wilkins et al pooled return to work data from four studies and also found no statistical significant difference.⁵ Soroceanu et al reported a statistically significant difference with the pooled data from four studies; operatively treated patients returned to work 19 days earlier than nonoperatively treated patients ($P=0.0014$).⁶ Wilkins et al included return to work data in their pooled results from the studies by Nistor et al and Cetti et al.^{5 52 57} In our study, we did not use the return to work data from these two studies owing to reporting of mean and range and the absence of standard deviations. Soroceanu et al also included the study by Cetti et al, as well as the study by Majewski et al,^{6 52 62} which we excluded

Table 3 | Secondary sensitivity analyses of studies included in meta-analysis of Achilles tendon ruptures

Studies	Re-rupture					Complication				
	No	RD (%)	RR (95% CI)	P value	I ² (%)	No	RD (%)	RR (95% CI)	P value	I ² (%)
All studies	29	1.6	0.43 (0.31 to 0.60)	<0.001	22	26	3.3	2.76 (1.84 to 4.13)	<0.001	45
High quality studies	17	5.1	0.44 (0.30 to 0.64)	<0.001	0	16	8.8	2.72 (1.44 to 5.12)	0.002	62
Study period (2000 or after)	14	0.9	0.59 (0.42 to 0.83)	0.002	10	14	2.4	2.15 (1.28 to 3.60)	0.004	52

RD=risk difference; RR=risk ratio.

as it was in a language other than English. However, both our meta-analysis and the studies by Wilkins et al and Soroceanu et al are limited by the number of included patients in the return to work subgroup analyses.^{5 6} Unfortunately, accurate comparison of functional outcome measures remains difficult owing to differences in protocols, patient oriented outcome measures, duration of follow-up, and presentation of data.

We found a lower re-rupture rate after both early and late full weight bearing in favour of operative treatment; this is in contrast to a previous meta-analysis by Van der Eng et al,⁶³ which found no difference in re-rupture rate. The previous meta-analysis could be limited by the number of included patients in the subgroup analyses. In our review, with the addition of observational studies, sensitivity analysis showed a significant difference in re-rupture rate after both early and late full weight bearing in favour of operative treatment. However, regardless of re-rupture rate, timing of weight bearing might influence other outcome measures as shown in different lower extremity injuries. De Boer et al found that early weight bearing regimens did not negatively affect functional outcome after treatment for displaced intra-articular calcaneal fractures.⁶⁴ Previously, Smeeing et al showed that early weight bearing tended to accelerate return to work and daily activities compared with late weight bearing, after internal fixation of ankle fractures.⁶⁵ Eliasson et al evaluated tendon elongation, mechanical properties, and functional outcomes during the first 12 months after operative treatment of acute Achilles tendon ruptures.⁶⁶ However, they found that different rehabilitation regimens did not affect the outcome measures. Further research could focus on the effect of early weight bearing and long term functional outcome after treatment of Achilles tendon ruptures.

Soroceanu et al found no significant difference in re-rupture rate in their subgroup analysis if functional rehabilitation with early range of motion was used (risk difference 1.7%; $P=0.45$).⁶ However, they did not define the specific inclusion criteria and definition of early range of motion and functional rehabilitation. Unfortunately, evaluation of the effect of accelerated functional rehabilitation remains difficult owing to use of a wide variety of definitions and protocols. Our review found no significant difference in re-rupture rate if accelerated functional rehabilitation with early range of motion within three weeks was used after nonoperative treatment. These findings indicate that nonoperative management is acceptable for acute Achilles tendon ruptures, if patients are instructed and monitored according to a standardised rehabilitation protocol. However, both our review and the study by Soroceanu et al could be limited by the number of included patients in the subgroup analyses.⁶

The sensitivity analyses including high quality studies resulted in similar risk ratios and significance levels for re-rupture and complication rate. The results showed a risk difference of 5.1% for re-rupture rate, comparable to previous results of meta-analyses of

RCTs alone. However, the risk difference of other complications (8.8%) in the high quality sensitivity analysis was still considerably lower than in previous reports. This difference in other complications could be attributable to the inclusion of studies with both open and minimally invasive surgical techniques. The complication sensitivity analysis with high quality studies included one RCT with minimally invasive surgery and three observational studies that included both open and minimally invasive surgery. A meta-analysis by Yang et al,⁶⁷ including five RCTs and four cohort studies, found a significantly lower rate of deep infection with percutaneous treatment (0.6%) than with open treatment (3.6%) ($P=0.04$). However, the authors reported no significant difference in the rate of re-rupture between percutaneous and open treatment.⁶⁷

The sensitivity analyses including studies with a study period after the year 2000 showed similar risk ratios and significance levels regarding re-rupture and complication rate. However, the risk differences between treatment groups were smaller than in all other analyses. The study period sensitivity analyses included one RCT with minimally invasive surgery and four observational studies that included both open and minimally invasive surgery. These findings might indicate an overall reduction in complications after treatment of Achilles tendon ruptures due to the development of new rehabilitation protocols and operative techniques, regardless the use of operative or nonoperative treatment. However, it should be noted that both the level of high quality studies and the study period were arbitrarily chosen.

We found no difference in pooled effect estimates from RCTs and observational studies. This is in line with previous reports showing that differences in effect estimates between RCTs and observational studies are small.^{8 9 11 13 15 17} Observational studies, however, have also been associated with an overestimation of treatment effects compared with RCTs.^{68 69} Hemkens et al assessed the difference in treatment effect estimates for mortality between observational studies and RCTs.⁶⁹ They evaluated 16 observational studies and 36 subsequent RCTs investigating the same clinical questions. Overall, observational studies significantly overestimated the effects of treatment compared with RCTs.⁶⁹ This overestimation of treatment effects could be explained by the effects of bias and confounding in observational studies.⁷⁰ However, overestimates by observational studies could also be explained by the potential selection bias in RCTs. RCTs require strict conditions such as selection of participants, inclusion/exclusion criteria, randomisation method, and outcome measurements. The patient population in daily clinical practice can differ from the often highly selected patient populations in RCTs, which could be the reason for the discrepancy between treatment effects.^{71 72} Nevertheless, observational studies increase sample size, which could lead to the evaluation of small treatment effects and infrequent outcome measures. Furthermore, the addition of observational studies might provide insight into a

variety of populations and long term effects. These results could improve the representation of daily clinical practice, with various levels of surgical experience and differences in operative techniques, provided that confounding has been adequately addressed.^{12 13} In this meta-analysis, pooled effect estimates obtained from RCTs and observational studies were similar. Several orthopaedic trauma meta-analyses including both RCTs and observational studies have shown high quality observational studies to result in similar treatment effects to RCTs.¹⁵⁻¹⁷ These findings indicate that the effect of potentially unmeasured confounding in high quality observational studies seems relatively small, emphasising the possible benefits of combining different study designs for the evaluation of objective outcome measures after surgical treatment.

Limitations of this study

Several potential limitations in this review need to be considered. Firstly, results might be influenced by missing articles. However, in addition to the extensive electronic database search, funnel plots did not indicate evidence for publication bias. Three studies could not be obtained in full text, but these articles were all published before 1996.²⁶⁻²⁸ Secondly, the methodological quality of included studies was assessed by the MINORS criteria, which do not differentiate between randomised and non-randomised studies. However, the MINORS criteria were externally validated using RCTs and are able to distinguish adequately between study designs, as well designed randomised trials score higher than well designed non-randomised studies.²¹ The incidence of complications could be affected by the use of different treatment protocols. Five studies mentioned the use of prophylaxis for deep vein thrombosis.^{51 53 56 58 59} However, descriptions were not comprehensive and the duration and types of prophylaxis varied widely. Finally, sensitivity analyses for the evaluation of weight bearing status and accelerated rehabilitation were performed using data from both RCTs and observational studies. However, the primary analysis showed no significant difference in effect estimates between the two study designs in terms of re-rupture rate.

Implications for future research

Operative treatment of acute Achilles tendon ruptures reduces the risk of re-rupture compared with nonoperative treatment, although the incidence of re-ruptures is low and differences are small (2.3% v 3.9%). Operative treatment results in a higher risk of other complications compared with nonoperative treatment, mostly attributable to the increased risk of infection. Nonoperative treatment might be the preferred treatment for acute Achilles tendon rupture, owing to the higher risk of other complications after operative treatment and the relative small benefit in re-rupture rate. However, patient specific factors should always be taken into consideration and patients should be counselled about the incidence of complications.

Unfortunately, comparison of the literature remains difficult owing to a wide variety of rehabilitation

protocols, weight bearing restrictions, treatment modalities, patient oriented outcome measures, and duration of follow-up. The discordance among studies makes comparisons between treatment modalities difficult, indicating a substantial need for further research. We suggest future research to focus on the effect of comorbidities on the success of treatment for Achilles tendon rupture. Studies could compare outcomes according to different age groups and evaluate effects in a variety of populations such as in patients with immunosuppression, diabetes mellitus, increased body mass index, neuropathy, peripheral vascular disease, and dermatological disorders. Furthermore, future studies should strive to determine the optimal treatment for acute Achilles tendon ruptures on the basis of patients' expectations. Operative treatment is associated with complications inherent to the treatment itself, such as infection. However, athletic people may prefer operative treatment to enhance and expedite their outcomes, whereas a sedentary person with limited functional outcome expectations may prefer nonoperative treatment.³ We believe that more data are needed for the development of a shared decision making algorithm to guide surgeons and physicians regarding the most appropriate treatment option for each individual patient.

Conclusions

In this meta-analysis, operative treatment of acute Achilles tendon ruptures reduced the risk of re-rupture compared with nonoperative treatment. However, re-rupture rates are low and differences between treatment groups are small, with a risk difference of 1.6%. Operative treatment results in a higher risk of other complications, with a risk difference of 3.3%, mostly due to the increased risk of infection. Patients should be counselled about complications, and the final decision for operative or nonoperative management should be based on patient specific factors and shared decision making. Further research is needed for the development of a shared decision making algorithm. Moreover, this review emphasises the potential benefits of adding high quality observational studies in meta-analyses to complement RCTs for the evaluation of objective outcome measures after surgical treatments.

AUTHOR AFFILIATIONS

¹Department of Surgery, University Medical Center Utrecht, 3508 GA Utrecht, Netherlands

²Department of Orthopedic Surgery, Harvard Medical School Orthopedic Trauma Initiative, Massachusetts General Hospital, Boston, MA, USA

³Department of Surgery, Diaconessenhuis Hospital, Utrecht, Netherlands

⁴Department of Surgery, Academic Medical Center, Amsterdam, Netherlands

⁵Department of Surgery, St. Antonius Hospital, Nieuwegein, Netherlands

⁶Department of Orthopedic Surgery, Spaarne Hospital, Hoofddorp, Netherlands

⁷Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, Netherlands

Contributors: RMH and RHHG conceived and designed the study. YO, RHHG, RBB, and RMH acquired, analysed, and interpreted the data. YO drafted the first version of the manuscript. YO, OvdM, RHHG, RBB, MH, and RMH drafted the subsequent versions of the manuscript. MVH, FH, LPHL, and DvdV gave clinical advice in interpreting the results. All authors contributed critically to subsequent revisions and approved the final manuscript. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. YO is the guarantor.

Funding: None.

Competing interests: All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Ethical approval: Not needed.

Data sharing: No additional data available.

Transparency: The lead author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

- 1 Ganestam A, Kallelose T, Troelsen A, Barfod KW. Increasing incidence of acute Achilles tendon rupture and a noticeable decline in surgical treatment from 1994 to 2013. A nationwide registry study of 33,160 patients. *Knee Surg Sports Traumatol Arthrosc* 2016;24:3730-7. doi:10.1007/s00167-015-3544-5
- 2 Egger AC, Berkowitz MJ. Achilles tendon injuries. *Curr Rev Musculoskelet Med* 2017;10:72-80. doi:10.1007/s12178-017-9386-7
- 3 Khan RJ, Carey Smith RL. Surgical interventions for treating acute Achilles tendon ruptures. *Cochrane Database Syst Rev* 2010;(9):CD003674.
- 4 Jiang N, Wang B, Chen A, Dong F, Yu B. Operative versus nonoperative treatment for acute Achilles tendon rupture: a meta-analysis based on current evidence. *Int Orthop* 2012;36:765-73. doi:10.1007/s00264-011-1431-3
- 5 Wilkins R, Bisson LJ. Operative versus nonoperative management of acute Achilles tendon ruptures: a quantitative systematic review of randomized controlled trials. *Am J Sports Med* 2012;40:2154-60. doi:10.1177/0363546512453293
- 6 Soroceanu A, Sidhwa F, Aarabi S, Kaufman A, Glazebrook M. Surgical versus nonsurgical treatment of acute Achilles tendon rupture: a meta-analysis of randomized trials. *J Bone Joint Surg Am* 2012;94:2136-43. doi:10.2106/JBJS.K.00917
- 7 Erickson BJ, Mascarenhas R, Saltzman BM, et al. Is Operative Treatment of Achilles Tendon Ruptures Superior to Nonoperative Treatment?: A Systematic Review of Overlapping Meta-analyses. *Orthop J Sports Med* 2015;3:2325967115579188. doi:10.1177/2325967115579188
- 8 Benson K, Hartz AJ. A comparison of observational studies and randomized, controlled trials. *N Engl J Med* 2000;342:1878-86. doi:10.1056/NEJM200006223422506
- 9 Concato J, Shah N, Horwitz RJ. Randomized, controlled trials, observational studies, and the hierarchy of research designs. *N Engl J Med* 2000;342:1887-92. doi:10.1056/NEJM200006223422507
- 10 Anglemeyer A, Horvath HT, Bero L. Healthcare outcomes assessed with observational study designs compared with those assessed in randomized trials. *Cochrane Database Syst Rev* 2014;(4):MR000034.
- 11 Abraham NS, Byrne CJ, Young JM, Solomon MJ. Meta-analysis of well-designed nonrandomized comparative studies of surgical procedures is as good as randomized controlled trials. *J Clin Epidemiol* 2010;63:238-45. doi:10.1016/j.jclinepi.2009.04.005
- 12 Arditi C, Burnand B, Poytremann-Bridevaux I. Adding non-randomised studies to a Cochrane review brings complementary information for healthcare stakeholders: an augmented systematic review and meta-analysis. *BMC Health Serv Res* 2016;16:598. doi:10.1186/s12913-016-1816-5
- 13 Frieden TR. Evidence for Health Decision Making - Beyond Randomized, Controlled Trials. *N Engl J Med* 2017;377:465-75. doi:10.1056/NEJMra1614394
- 14 Houwert RM, Smeeing DP, Ahmed Ali U, Hietbrink F, Kruyt MC, van der Meijden OA. Plate fixation or intramedullary fixation for midshaft clavicle fractures: a systematic review and meta-analysis of randomized controlled trials and observational studies. *J Shoulder Elbow Surg* 2016;25:1195-203. doi:10.1016/j.jse.2016.01.018
- 15 Smeeing DP, van der Ven DJC, Hietbrink F, et al. Surgical Versus Nonsurgical Treatment for Midshaft Clavicle Fractures in Patients Aged 16 Years and Older: A Systematic Review, Meta-analysis, and Comparison of Randomized Controlled Trials and Observational Studies. *Am J Sports Med* 2017;45:1937-45. doi:10.1177/0363546516673615
- 16 Beks RB, Ochen Y, Frima H, et al. Operative versus nonoperative treatment of proximal humeral fractures: a systematic review, meta-analysis, and comparison of observational studies and randomized controlled trials. *J Shoulder Elbow Surg* 2018;27:1526-34. doi:10.1016/j.jse.2018.03.009
- 17 Beks RB, Peek J, de Jong MB, et al. Fixation of flail chest or multiple rib fractures: current evidence and how to proceed. A systematic review and meta-analysis. *Eur J Trauma Emerg Surg* 2018. doi:10.1007/s00068-018-1020-x
- 18 Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008-12. doi:10.1001/jama.283.15.2008
- 19 Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006-12. doi:10.1016/j.jclinepi.2009.06.005
- 20 Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 2009;62:e1-34. doi:10.1016/j.jclinepi.2009.06.006
- 21 Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ J Surg* 2003;73:712-6. doi:10.1046/j.1445-2197.2003.02748.x
- 22 Nilsson-Helander K, Thomeé R, Silbernagel KG, et al. The Achilles tendon Total Rupture Score (ATRS): development and validation. *Am J Sports Med* 2007;35:421-6. doi:10.1177/0363546506294856
- 23 Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). 2011. <http://www.cochrane.org/training/cochrane-handbook>.
- 24 Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-34. doi:10.1136/bmj.315.7109.629
- 25 Review Manager (RevMan) [Computer program]. Version 5.3. The Nordic Cochrane Centre, The Cochrane Collaboration, 2014
- 26 Lill H, Moor C, Fecht E, et al. Achilles tendon rupture - Operative or conservative functional treatment? *Aktuelle Traumatol* 1996;26:95-100.
- 27 Häggmark T, Liedberg H, Eriksson E, Wredmark T. Calf muscle atrophy and muscle function after non-operative vs operative treatment of achilles tendon ruptures. *Orthopedics* 1986;9:160-4.
- 28 Inglis AE, Scott WN, Sculco TP, Patterson AH. Ruptures of the tendo achillis. An objective assessment of surgical and non-surgical treatment. *J Bone Joint Surg Am* 1976;58:990-3. doi:10.2106/00004623-197658070-00015
- 29 Möller M, Lind K, Movin T, Karlsson J. Calf muscle function after Achilles tendon rupture. A prospective, randomised study comparing surgical and non-surgical treatment. *Scand J Med Sci Sports* 2002;12:9-16. doi:10.1034/j.1600-0838.2002.120103.x
- 30 Olsson N, Nilsson-Helander K, Karlsson J, et al. Major functional deficits persist 2 years after acute Achilles tendon rupture. *Knee Surg Sports Traumatol Arthrosc* 2011;19:1385-93. doi:10.1007/s00167-011-1511-3
- 31 Olsson N, Karlsson J, Eriksson B, Brorsson A, Lundberg M, Silbernagel KG. Ability to perform a single heel-rise is significantly related to patient-reported outcome after Achilles tendon rupture. *Scand J Med Sci Sports* 2014;24:152-8. doi:10.1111/j.1600-0838.2012.01497.x
- 32 Westin O, Nilsson Helander K, Grävare Silbernagel K, Möller M, Kälebo P, Karlsson J. Acute Ultrasonography Investigation to Predict Reruptures and Outcomes in Patients With an Achilles Tendon Rupture. *Orthop J Sports Med* 2016;4:2325967116667920. doi:10.1177/2325967116667920
- 33 Costa ML, MacMillan K, Halliday D, et al. Randomised controlled trials of immediate weight-bearing mobilisation for rupture of the tendo Achillis. *J Bone Joint Surg Br* 2006;88:69-77. doi:10.1302/0301-620X.88B1.16549
- 34 Jackson G, Sinclair VF, McLaughlin C, Barrie J. Outcomes of functional weight-bearing rehabilitation of Achilles tendon ruptures. *Orthopedics* 2013;36:e1053-9. doi:10.3928/01477447-20130724-23
- 35 Kotnis R, David S, Handley R, Willett K, Ostlere S. Dynamic ultrasound as a selection tool for reducing achilles

- tendon reruptures. *Am J Sports Med* 2006;34:1395-400. doi:10.1177/0363546506288678
- 36 Bergkvist D, Åström I, Josefsson P-O, Dahlberg LE. Acute Achilles tendon rupture: a questionnaire follow-up of 487 patients. *J Bone Joint Surg Am* 2012;94:1229-33. doi:10.2106/JBJS.J.01601
- 37 Carden DG, Noble J, Chalmers J, Lunn P, Ellis J. Rupture of the calcaneal tendon. The early and late management. *J Bone Joint Surg Br* 1987;69:416-20. doi:10.1302/0301-620X.69B3.3294839
- 38 Cukelj F, Bandalovic A, Knezevic J, Pavic A, Pivalica B, Bakota B. Treatment of ruptured Achilles tendon: Operative or non-operative procedure? *Injury* 2015;46(Suppl 6):S137-42. doi:10.1016/j.injury.2015.10.070
- 39 Ebinesan AD, Sarai BS, Walley GD, Maffulli N. Conservative, open or percutaneous repair for acute rupture of the Achilles tendon. *Disabil Rehabil* 2008;30:1721-5. doi:10.1080/09638280701786815
- 40 Fahlström M, Björnstig U, Lorentzon R. Acute Achilles tendon rupture in badminton players. *Am J Sports Med* 1998;26:467-70. doi:10.1177/03635465980260032201
- 41 Grubor P, Grubor M. Treatment of Achilles tendon rupture using different methods. *Vojnosanit Pregl* 2012;69:663-8. doi:10.2298/VSP1208663G
- 42 Gwynne-Jones DP, Sims M, Handcock D. Epidemiology and outcomes of acute Achilles tendon rupture with operative or nonoperative treatment using an identical functional bracing protocol. *Foot Ankle Int* 2011;32:337-43. doi:10.3113/FAI.2011.0337
- 43 Jaakkola JI, Beskin JL, Griffith LH, Cernansky G. Early ankle motion after triple bundle technique repair vs. casting for acute Achilles tendon rupture. *Foot Ankle Int* 2001;22:979-84. doi:10.1177/107110070102201210
- 44 Lim CS, Lees D, Gwynne-Jones DP. Functional Outcome of Acute Achilles Tendon Rupture With and Without Operative Treatment Using Identical Functional Bracing Protocol. *Foot Ankle Int* 2017;38:1331-6. doi:10.1177/1071100717728687
- 45 Miller D, Waterston S, Reaper J, Barrass V, Maffulli N. Conservative management, percutaneous or open repair of acute Achilles tendon rupture: a retrospective study. *Scott Med J* 2005;50:160-5. doi:10.1177/003693300505000408
- 46 Nestorson J, Movin T, Möller M, Karlsson J. Function after Achilles tendon rupture in the elderly: 25 patients older than 65 years followed for 3 years. *Acta Orthop Scand* 2000;71:64-8. doi:10.1080/00016470052943928
- 47 Rajasekar K, Gholve P, Faraj AA, Kosygan KP. A subjective outcome analysis of tendo-Achilles rupture. *J Foot Ankle Surg* 2005;44:32-6. doi:10.1053/j.jfas.2004.11.008
- 48 Renninger CH, Kuhn K, Fellars T, Youngblood S, Bellamy J. Operative and Nonoperative Management of Achilles Tendon Ruptures in Active Duty Military Population. *Foot Ankle Int* 2016;37:269-73. doi:10.1177/1071100715615322
- 49 van der Linden-van der Zwaag HMJ, Nelissen RGH, Sintenie JB. Results of surgical versus non-surgical treatment of Achilles tendon rupture. *Int Orthop* 2004;28:370-3. doi:10.1007/s00264-004-0575-9
- 50 Wang D, Sandlin MI, Cohen JR, Lord EL, Petrigliano FA, SooHoo NF. Operative versus nonoperative treatment of acute Achilles tendon rupture: An analysis of 12,570 patients in a large healthcare database. *Foot Ankle Surg* 2015;21:250-3. doi:10.1016/j.jfas.2015.01.009
- 51 Weber M, Niemann M, Lanz R, Müller T. Nonoperative treatment of acute rupture of the achilles tendon: results of a new protocol and comparison with operative treatment. *Am J Sports Med* 2003;31:685-91. doi:10.1177/03635465030310050901
- 52 Cetti R, Christensen SE, Ejsted R, Jensen NM, Jorgensen U. Operative versus nonoperative treatment of Achilles tendon rupture. A prospective randomized study and review of the literature. *Am J Sports Med* 1993;21:791-9. doi:10.1177/036354659302100606
- 53 Keating JF, Will EM. Operative versus non-operative treatment of acute rupture of tendo Achillis: a prospective randomised evaluation of functional outcome. *J Bone Joint Surg Br* 2011;93:1071-8. doi:10.1302/0301-620X.93B8.25998
- 54 Lantto I, Heikkinen J, Flinkkila T, et al. A Prospective Randomized Trial Comparing Surgical and Nonsurgical Treatments of Acute Achilles Tendon Ruptures. *Am J Sports Med* 2016;44:2406-14. doi:10.1177/0363546516651060
- 55 Metz R, Verleisdonk EJ, van der Heijden GJ, et al. Acute Achilles tendon rupture: minimally invasive surgery versus nonoperative treatment with immediate full weightbearing--a randomized controlled trial. *Am J Sports Med* 2008;36:1688-94. doi:10.1177/0363546508319312
- 56 Möller M, Movin T, Granhed H, Lind K, Faxén E, Karlsson J. Acute rupture of tendon Achillis. A prospective randomised study of comparison between surgical and non-surgical treatment. *J Bone Joint Surg Br* 2001;83:843-8. doi:10.1302/0301-620X.83B6.0830843
- 57 Nistor L. Surgical and non-surgical treatment of Achilles Tendon rupture. A prospective randomized study. *J Bone Joint Surg Am* 1981;63:394-9. doi:10.2106/00004623-198163030-00012
- 58 Nilsson-Helander K, Silbernagel KG, Thomeé R, et al. Acute achilles tendon rupture: a randomized, controlled study comparing surgical and nonsurgical treatments using validated outcome measures. *Am J Sports Med* 2010;38:2186-93. doi:10.1177/0363546510376052
- 59 Olsson N, Silbernagel KG, Eriksson BI, et al. Stable surgical repair with accelerated rehabilitation versus nonsurgical treatment for acute Achilles tendon ruptures: a randomized controlled study. *Am J Sports Med* 2013;41:2867-76. doi:10.1177/0363546513503282
- 60 Twaddle BC, Poon P. Early motion for Achilles tendon ruptures: is surgery important? A randomized, prospective study. *Am J Sports Med* 2007;35:2033-8. doi:10.1177/0363546507307503
- 61 Willits K, Amendola A, Bryant D, et al. Operative versus nonoperative treatment of acute Achilles tendon ruptures: a multicenter randomized trial using accelerated functional rehabilitation. *J Bone Joint Surg Am* 2010;92:2767-75. doi:10.2106/JBJS.101401
- 62 Majewski M, Rickert M, Steinbrück K. [Achilles tendon rupture. A prospective study assessing various treatment possibilities]. *Orthopade* 2000;29:670-6.
- 63 van der Eng DM, Schepers T, Goslings JC, Schep NW. Rerupture rate after early weightbearing in operative versus conservative treatment of Achilles tendon ruptures: a meta-analysis. *J Foot Ankle Surg* 2013;52:622-8. doi:10.1053/j.jfas.2013.03.027
- 64 De Boer AS, Van Lieshout EMM, Van Moolenbroek G, Den Hartog D, Verhofstad MHJ. The effect of time to post-operative weightbearing on functional and clinical outcomes in adults with a displaced intra-articular calcaneal fracture; A systematic review and pooled analysis. *Injury* 2018;49:743-52. doi:10.1016/j.injury.2018.02.021
- 65 Smeeing DJ, Houwert RM, Briet JP, et al. Weight-bearing and mobilization in the postoperative care of ankle fractures: a systematic review and meta-analysis of randomized controlled trials and cohort studies. *PLoS One* 2015;10:e0118320. doi:10.1371/journal.pone.0118320
- 66 Eliasson P, Agergaard A-S, Couppe C, et al. The Ruptured Achilles Tendon Elongates for 6 Months After Surgical Repair Regardless of Early or Late Weightbearing in Combination With Ankle Mobilization: A Randomized Clinical Trial. *Am J Sports Med* 2018;46:2492-502. doi:10.1177/0363546518781826
- 67 Yang B, Liu Y, Kan S, et al. Outcomes and complications of percutaneous versus open repair of acute Achilles tendon rupture: A meta-analysis. *Int J Surg* 2017;40:178-86. doi:10.1016/j.ijsu.2017.03.021
- 68 Ioannidis JP, Haidich AB, Pappa M, et al. Comparison of evidence of treatment effects in randomized and nonrandomized studies. *JAMA* 2001;286:821-30. doi:10.1001/jama.286.7.821
- 69 Hemkens LG, Contopoulos-Ioannidis DG, Ioannidis JPA. Agreement of treatment effects for mortality from routinely collected data and subsequent randomized trials: meta-epidemiological survey. *BMJ* 2016;352:i493. doi:10.1136/bmj.i493
- 70 Boyko EJ. Observational research--opportunities and limitations. *J Diabetes Complications* 2013;27:642-8. doi:10.1016/j.jdiacomp.2013.07.007
- 71 Van Spall HGC, Toren A, Kiss A, Fowler RA. Eligibility criteria of randomized controlled trials published in high-impact general medical journals: a systematic sampling review. *JAMA* 2007;297:1233-40. doi:10.1001/jama.297.11.1233
- 72 Khan AY, Preskorn SH, Baker B. Effect of study criteria on recruitment and generalizability of the results. *J Clin Psychopharmacol* 2005;25:271-5. doi:10.1097/01.jcp.0000161497.73514.80

Supplementary materials